UNCLASSIFIED

AD NUMBER AD064663 CLASSIFICATION CHANGES TO: UNCLASSIFIED FROM: CONFIDENTIAL LIMITATION CHANGES

TO:

Approved for public release; distribution is unlimited.

FROM:

Distribution authorized to DoD only; Administrative/Operational Use; 20 DEC 1951. Other requests shall be referred to Naval Ordnance Laboratory, Attn: Explosives Research Department, White Oak, MD. Pre-dates formal DoD distribution statements. Treat as DoD only.

AUTHORITY

31 Dec 1963, DoDD 5200.10; USNSWC ltr dtd 4 Dec 1974

UNCLASSIFIED

AD _____

Reproduced by the

ARMED SERVICES TECHNICAL INFORMATION AGENCY ARLINGTON HALL STATION ARLINGTON 12, VIRGINIA



DOWNGRADED AT 3 YEAR INTERVALS: DECLASSIFIED AFTER 12 YEARS DOD DIR 5200.10

UNCLASSIFIED

ADG AGG G

Armed Services Technical Information Agency

Reproduced by DOCUMENT SERVICE CENTER KNOTT BUILDING, DAYTON, 2, 0 HIO

Because of our limited supply, you are requested to RETURN THIS COPY WHEN IT HAS SERVED YOUR PURPOSE so that it may be made available to other requesters.

Your cooperation will be appreciated.

NOTICE: WHEN GOVERNMENT OR OTHER DRAWINGS, SPECIFICATIONS OR OTHER DATA ARE USED FOR ANY PURPOSE OTHER THAN IN CONNECTION WITH A DEFINITELY RELATED GOVERNMENT PROCUREMENT OPERATION, THE U. S. GOVERNMENT THEREBY INCURS NO RESPONSIBILITY, NOR ANY OBLIGATION WHATSOEVER; AND THE FACT THAT THE GOVERNMENT MAY HAVE FORMULATED, FURNISHED, OR IN ANY WAY SUPPLIED THE SAID DRAWINGS, SPECIFICATIONS, OR OTHER DATA IS NOT TO BE REGARDED BY IMPLICATION OR OTHERWISE AS IN ANY MANNER LICENSING THE HOLDER OR ANY OTHER PERSON OR CORPORATION, OR CONVEYING ANY RIGHTS OR PERMISSION TO MANUFACTURE, USE OR SELL ANY PATENTED INVENTION THAT MAY IN ANY WAY BE RELATED THERETO.

64663 FILE COPY

NAVORD REPORT 2307



PRIMARY EXPLOSIVES RESEARCH

III. Synthesis and Examination of 5-Amino-1,2,3,4-Thiatriazole and the Metal Salts of 3,5-Dinitro-1,2,4-Triazole

20 December 1951



U. S. NAVAL ORDNANCE LABORATORY
WHITE OAK, MARYLAND



SECURITY INFORMATION

This document contains information affecting the national defense of the United States within the meaning of the Espionage Laws, Title 18, U.S.C., Sections 793 and 794, the transmission or revelation of which in any manner to an unauthorized person, is prohibited by law.

Reproduction of this document in any form by other than activities of the Department of Defense and the Atomic Energy Commission is not authorized unless specifically approved by the Secretary of the Navy or the Chief of Naval Operations as appropriate.



PRIMARY EXPLOSIVES RESEARCH

III. Synthesis and Examination of 5-Amino-1,2,3,4-Thiatriazole and the Metal Salts of 3,5-Dinitro-1,2,4-Triazole

Prepared by:

FRANCIS TAYLOR, JR.

Approved by: Darrell V. Sickman Chief, Chemistry Division

ABSTRACT: The following metal salts of 3,5-dinitro-1,2,4-triazole were prepared and examined as possible new primary explosives.

Gupric Lead Silver

Ferric Mercuric Tin

Ferrous Mercurous Zinc

All but the silver and mercurous salts were rejected because of excessive water solubility. These compounds had satisfactory thermal stability, but their impact sensitivities were in the range of that of RDX. Further tests are being made on the silver salt and the results will be reported elsewhere.

5-Amino-1,2,3,4-thiatriazole was prepared and examined as a possible primary emplosive. Its impact sensitivity was in the range of that of lead azide, but the compound was thermally unstable. No further work on it is contemplated.

Explosives Research Department U. S. NAVAL ORDNANCE LABORATORY White Oak, Maryland

CONFIDENTIAL



CONFIDENTIAL

NAVORD Report 2307

20 December 1951

This report, for information only, is a description of the preparation of the heavy metal salts, particularly the silver and mercurous, of 3,5-dinitro-1,2,4-triazole, and of 5-amino-1,2,3,4-thiatriazole, performed under Task NOL-Re2b-41-1-52. The reliability of the work and the validity of the conclusions are the responsibility of the author and of the Chemistry Division, Explosives Research Department, of the U. S. Naval Ordnance Laboratory.

W. G. SCHINDLER Rear Admiral, USN Commander

J. E. ABLARD, Deputy Chief Explosives Research Department

By direction

11 CONFIDENTIAL

•	Tab	ole d	of Co	onte	nts	•					
	********				eri,tar.	'				Pa	ge No.
Introduction											1
Experimental					•		•	٠	•	•	ī
Silver 3,5-dinitro											ī
Mercurous 3,5-dini	tro-1	1,2,1	l-tr	lazo	le		•	•	•	•	2
5-Amino-1,2,3,4-th											3
Physical Properties,	Tab1	Le I	• •		•			•	•	•	4
Conclusions					•			•	•		5
Acknowledgements										•	5
References											ź.

PRIMARY EXPLOSIVES RESEARCH

III. Synthesis and Examination of 5-Amino-1,2,3,4-Thiatriazole and the Metal Salts of 3,5-Dinitro-1,2,4-Triazole

INTRODUCTION

- 1. Working on a Bureau of Ordnance Contract for high explosives research, the Naugatuck Chemical Division of the United States Rubber Company discovered a method of preparing 3,5-dinitro-1,2,4-triazole and prepared several of its salts. They suggested that the silver and lead salts might find some use as primary explosives, since they were too sensitive for consideration as high explosives.
- 2. Of the following metal salts, the only ones insoluble in water were the silver and mercurous compounds.

Cupric Lead Silver

Ferric Mercuric Stannous

Ferrous Mercurous Zinc

It is believed that water insolubility is a desirable physical property for a compound being considered as a possible new primary explosive. Therefore all of the water soluble salts were dropped from consideration.

3. Freund and Schander reported the preparation of 5-amino-1,2,3,4-thiatriazole in the early literature (2). The Naugatuck Chemical Division repeated the work and suggested the compound as a possible primary explosive (3).

EXPERIMENTAL

Silver Salt of 3,5-Dinitro-1,2,4-Triazole

4. The Naugatuck Chemical Division supplied a sample of the sodium salt of 3,5-dinitro-1,2,4-triazole, the starting material for the preparation of the silver and mercurous salts, prepared as described in their report (1). The sodium salt (2 grams, 0.01 mole) was dissolved in 20 cc of distilled water and stirred mechanically. To the aqueous sodium salt solution was added dropwise an aqueous solution of silver

CONFIDENTIAL

nitrate (1.8 grams, 0.01 moles in 15 cc of distilled water). The silver salt precipitated immediately. The reaction mixture was stirred for an additional ten minutes and then filtered on a Buchner funnel. The yellow solid was washed with liberal quantities of distilled water and finally dried under vacuum at 50°C. The salt was analyzed for silver by the method described by Pregl (4).

Calculated for C204N5Ag: Ag, 40.6

Found: Ag, 39.3, 41.27

The physical properties observed for the silver salt are tabulated in Table I.

Mercurous Salt of 3,5-Dinitro-1,2,4-Triazole

5. The mercurous salt of 3,5-dinitro-1,2,4-triazole was also prepared using the sodium salt as the starting material. An aqueous solution of mercurous nitrate was prepared by dissolving 15 grams (0.055 moles) in 500 cc of distilled water containing 5 cc of concentrated nitric acid. The aqueous mercurous nitrate was stirred mechanically while an aqueous solution of the sodium salt (4.5 grams, 0.025 moles, in 15 cc of distilled water) was added dropwise. On addition of the sodium salt a yellow solid precipitated immediately. The solid was filtered, washed with distilled water, and dried under vacuum at 50°C. The solid was analyzed for mercury by the Oakwold Laboratory, Alexandria, Virginia.

Calculated for C204N5HE: Hg. 56.0

Found: Hg, 53.64, 53.82

This compound was reported by the Cakwold Laboratory to be very volatile and subject to violent decomposition, so that the accuracy of the analysis is open to some question. The physical properties observed for the mercurous salt are listed in Table I.

5-Amino-1,2,3,4-Thiatriazole

6. The method described by Freund and Schander for the preparation of 5-amino-1,2,3,4-thiatriazole was repeated (2). The crude material isolated from the reaction melted at 120°C. One recrystallization from ether raised the melting point 128-129°C. It was observed that sometimes the sample would melt in a capillary but that occasionally it would detonate. The literature reported the melting point as a detonation temperature. The physical properties observed are tabulated in Table I.

TEBLE I

Ignition Temperature	286°c /1.	(A)	128&
Density	2,781	des (en mis es) des	1 P. T. S.
Vacuum Stability ee gas/48 hrs/100°C	5°09 ec	0.12 00	27°25
Impact Sensitivity 2.5 kg weight	26 em	38. cm	8.6 cm
Compound	Silver-3,5.Dinitro 1,2,4-Trizzole	Marcurous-3,5-Binitro- 1,2,4-Trierole	Paranto-1,2,3,4-Thisa

Jourd to detonate in two out of two trials with a hot wire energized by 1000,000 ergs d

Enceeded the capacity of the thermal stability equipment (30 de) both at $100^{0}\mathrm{c}$ and $90^{\circ}\mathrm{c}$. 9

CONCLUSIONS

7. The silver and mercurous salts of 3,5-dinitro1,2,4-triazole exhibit satisfactory thermal stability but
neither appeared to possess the necessary sensitivity for
a primary explosive. The 5-amino-1,2,3,4-thiatriazole
possessed the necessary sensitivity for a primary explosive,
but had very poor thermal stability. A sample of the silver
salt was prepared for further evaluation by the Explosives
Properties Division, Explosives Research Department, and
the Chemistry Division, Engineering Department, of the
U. S. Naval Ordnance Laboratory. These results will be
reported elsewhere. No further work is planned on the
3,5-dinitro-1,2,4-triazole molecule.

ACKNOWLEDGEMENTS

8. The author wishes to acknowledge the assistance of J. M. Rosen for the thermal stability measurements and of B. O. Wilkerson for the metal analyses, both of the Chemistry Division, and of G. Svadeba, Explosives Properties Division, who made the impact sensitivity measurements.

REFERENCES

- 1. Naugatuck Chemical Division, United States Rubber Company, Nord 10,121, Progress Report No. 3, Summary from December 1947 to January 1949, dtd February 1949.
- 2. Freund, M. and Schander, A., Ber. 29, 2502 (1896).
- 3. Naugatuck Chemical Division, United States Rubber Company, NOrd 10,121, Monthly Progress Report, April 1949.
- 4. Pregl, F., Revised and edited by Grant, J., "Quantitative Organic Microanalysis", 4th edition, 1946.